

Growing Vegetables from Seed



Matt Jones

Horticulture Extension Agent NC Cooperative Extension - Chatham County Center















Tonight's Webinar

- Why grow from seed?
- Seed and seedling biology
- Planting calendars
- Seed starting methods
- Evaluations and resources







What is Cooperative Extension?

World's largest **non-formal education** network

Established 1914 by the Smith-Lever Act



Practical, non-degree programs





What is Cooperative Extension?

A nationwide network of

- Educators
- Researchers
- Volunteers













Additional Resources

Sustainable Vegetable Gardening Resources

- Many excellent Extension resources
- Slides from previous classes:
- Soils
- Pests & Diseases
- Warm season crops
- Cool season crops







NC STATE EXTENSION

NC Extension Gardener Handbook

https://go.ncsu.edu/eg-handbook



Free Online!

Hard copy – UNC Press (\$60)









Growing Vegetables from Seed



Matt Jones

Horticulture Extension Agent NC Cooperative Extension - Chatham County Center







Why grow veggies from seed?

- More cultivars
- Earlier harvests
- Less expensive
- Pest and disease avoidance
- Only cure for Gardener-Associated Winter Depression Syndrome (GAWDS)




Cucurbit downy mildew arrives in early summer.

Plant early to reduce yield loss.




Seeds: Baby Plants in a Box





Seed Anatomy






Figure 22-6c
Raven Biology of Plants, Eighth Edition
© 2013 W. H. Freeman and Company

Figure 22-6a


Raven Biology of Plants, Eighth Edition

© 2013 W. H. Freeman and Company

Cotyledon: Embryonic leaves, first to emerge after germination



Seedling Anatomy



Garden bean



Factors Affecting Germination

- Seeds must be alive
- Need the right environment
 - Temperature
 - Moisture
 - Air (oxygen)
 - Darkness/light
- Germination rates decline over time



Leftover seeds can be stored in an airtight container in a cool place



Types of Vegetable Varieties

Open Pollinated

'Heirloom' varieties – can save own seed and varieties will come true to type

Hybrid

- Result of a cross between 2 or more parents.
- Saved seed do not breed true
- Usually more uniform, more vigorous, more disease resistant

F₁ Hybrid

- Specific type of hybrid first generation
- Usually much more expensive!

GMO (Genetically Modified Organism, aka molecular breeding)

- Specific genes for specific traits incorporated via molecular biology
- Very few veg. crops: Sugar beet, apple, tomato, potato
- http://www.isaaa.org/gmapprovaldatabase/default.asp



Slide: Charlotte Glen - NCSU



Seed Packets



Common name and latin name of plant

Number of plants per seed packet

When to sow outside or inside

Planting depth, seed and row spacing, days until plants emerege and thining recommendations.

Gentically Modified Organism labling

Sell by date



Planting Seasons

Cool season

- Plant July-Sept for fall crop
- Feb April for spring crop

Warm season

- Plant after average last spring frost date, April 15
- See "Central NC Planting Calendar" for specific dates











Average Last Frost Date






Average First Frost Date





Planting Seasons



Slide: Charlotte Glen NC State Extension




Planting Calendars

colder temperatures and some frost

- Use Extension planting calendars
- Do not rely on seed packets for regionally-accurate information!

 Refer to direct planting, not seed starting



tolerate frost and should only be planted outside when frost is no longer a threat.

https://go.ncsu.edu/veggiecalendar



Table 1. Garden planting calendar for vegetables, fruits, and herbs in Central North Carolina.

	Days to Harvest	Dietanas Detween Blants	Jan	Fe	b	Ma	ar	Ар	r	Ma	ıy	Ju	ın	J	ul	Α	ug	Se	р	Oct	Nov	Dec
Fruit, Herb, or Vegetable	(from seed unless otherwise noted)	Distance Between Plants (inches)	115	1	15	1	15	1 1	5	1	15	1	15	1	15	1	15	1	15	115	1 15	1 15
Artichokes, globe	T = 1 year	30					Τ	Т	Т													
Artichokes, Jerusalem*	Tu = 6–8 months	9–12					Tu	Tu 1	ū													
Arugula	40-50	6–9		S	S	S	S									S	S	S	S			
Asparagus	C = 2 years	18			С	С	С															
Basil	T = 14–35 S = 50–75	2–8								S,T	S,T	S,T	S,T	S,T	S,T							
Beans, lima/bush	65-80	6							S	S	S	S	S	S	S							
Beans, lima/pole	75–95	6							S	S	S	S			S							
Beans, snap/bush	50-55	2					S	S	S	S	S	S	S	S	S	S	S	S	S			
Beans, snap/pole	65–70	6						S	S	S	S	S	S	S	S	S	S	S	S			
Beets	55-60	2				S	S	S							S	S	S	S				
Broccoli	T = 70-80	18			Τ	Τ	Τ	Т								Т	Τ	Τ				
Brussels sprouts	T = 40–50 S = 90–100**	14–18												Τ	T	Τ	Т					
Cabbage	T = 63–75 S = 90–120**	12		Τ	Т	Τ	Τ	Т							Τ	Т	Т	Τ				
Cabbage, Chinese	T = 45–55 S = 75–85	12					S,T									S	S		Τ	Т		
	Days to Harvest (from seed unless	Distance Between Plants	Jan	Fe	b	Ma	ar	Ар	r	Ma	ıy	Ju	ın	J	ul	Α	ug	Se	р	Oct	Nov	Dec
Fruit, Herb, or Vegetable	otherwise noted)	(inches)	115	1	15	1	15	1 1	5	1	15	1	15	1	15	1	15	1	15	115	1 15	1 15



When to Start Seeds

- Growing time before transplant varies by crop
- Count backwards from recommended transplanting date in planting calendar

Crop	Weeks	Crop	Weeks		
Broccoli	6-7	Kale	4-6		
Brussels Sprouts	6-7	Leeks	10-12		
Cabbage	6-7	Lettuce	5-6		
Celery	10-12	Okra	2-3		
Collards	5-7	Onions	10-12		
Cucumber	2-3	Peppers	8-10		
Eggplant	8-10	Tomatoes	6-8		







Tolerance to Transplanting

Transplant Well (Start Indoors)

- Broccoli
- Brussels Sprouts
- Cabbage
- Cauliflower
- Celery
- Collards
- Cucumber
- Eggplant

- Kohlrabi
- Kale
- Leeks
- Lettuce
- Melons
- Onions
- Peppers
- Squash
- Tomatoes

Transplant Poorly (Direct Seed)

- Beans
- Beet
- Carrot
- Corn
- Parsnip

- Peas
- Radish
- Rutabaga
- Spinach
- Turnip



Some crops are best seeded directly in the garden




Containers for Seed Starting



Plastic Flats





Plastic 4 and 6 Cell Packs





Plug Trays



Containers for Seed Starting

Peat Pots







Remove top + bottom or entire pot before planting







Containers for Seed Starting

Recycled Materials

- Anything with a drainage hole
- Disinfect with 1:10 bleach solution for 5 min., rinse and dry









Growing Media

Don't use soil from the garden!

- Weed seeds
- Poor drainage kills roots
 - Lack of O₂
 - Pathogens

Instead, use soilless substrates a.k.a. seed starting/potting mixes



Healthy



Nope!



Growing Media

Linivarcity









Vermiculite Perlite

University)	
Shredded sphagnum peat moss	10 gallons
No. 2, 3, or 4 domestic or African vermiculite ^b (horticultural grade, dust screened)	10 gallons
Pulverized Limestone Dolomitic Lime for mixes with domestic vermiculite or Calcitic Lime for mixes with African vermiculite	1 1/4 cups or 3/4 cups
Superphosphate (20% P) or Triple superphosphate (46%)	1/2 cup or 1/4 cup

Simple Seed Starting Mix (Rutgers



Growing Media

Seed starting mixes have the finest particles for the smallest seeds

- More expen\$ive
- Most regular potting mixes adequate
- You can mix well-screened compost (20%) with soilless media for additional nutrients









How to Plant Seeds

Plant according to recommendations

- Seed packet or Extension literature
- Depth = 1.5-2x seed diameter





Lightly cover seeds and carefully firm media for good soil contact

Exception: no need to cover lettuce seeds



Seed Sowing Strategies I

Sow many seed in a flat or pot, transplant individual seedlings to pots or 4 packs, etc.

- Efficient use of space
- Transplanting can help strengthen seedlings
- Works well for small seed and slower growing vegetables
- Best method when need individual plants





Slide: Charlotte Glen – NCSU



Solanaceous Crops

Individual Transplants





Cruciferous crops, head lettuce



Transplanting to Cells or Pots

- Transplant when first set of true leaves appear
- Lift from beneath with label, pencil or dibber
- Hold by cotyledon or leaf, NOT stem!
- Can transplant up to cotyledon, especially if leggy
- Keep out of direct sunlight for a day, water well







Seed Sowing Strategies II

Sow 1-3 seeds in a pot/cell (peat pot, 4 or 6 pack) to grow until large enough to transplant into garden

- Best for large seed (squash, cucumbers)
- Or plants grown in clumps/groups (lettuce, parsley)











Growing in Containers Outdoors







Grow Food in Small Spaces

Flexibility & Accessibility

Avoid Soil Problems



Growing in Containers Outdoors

More frequent watering

More frequent fertilization

Don't use native soil









Thinning Seedlings



When in doubt...

DECAPITATE!







NC COOPERATIVE EXTENSION



Pre-moisten media



Wrung-out sponge

Watering

Newly planted seeds

- Water carefully don't let seeds float away!
- Cover container to maintain high humidity

After germination

- Keep soil moist, but not wet
- When slightly wilting
- If you cannot squeeze-out water from top half-inch of media







Damping-off Diseases

Favored by cool, wet conditions

- Young seedlings more vulnerable to infection
- Rhizoctonia, Fusarium, Pythium

Symptoms

- Germination failure
- Seeds or seedlings soft, mushy, discolored
- Stems thin, roots absent or stunted

Prevention

- Don't over water
- Make sure containers are clean
- Don't over-apply fertilizer
- Provide adequate light







Light Requirements

Outdoor/Natural Light

- Greenhouse?
- Windows not enough

Indoors

- LED grow lights (\$\$)
- Fluorescent bulbs
 - T-8 or T-12 shop lights
 - Cool + warm
 - Broad spectrum grow lights



Keep lights 1-4 inches from seedlings for 12-16 hours per day



'Leggy' Seedlings

- Aim, for short, stocky, transplants
- Legginess caused by inadequate light









Heat Improves Germination Rates

Seedling heat mats are ideal



Crop	Min. (°F)	Optimum (°F)	Max. (°F)
Bean	60	75-85	95
Broccoli	40	60-85	95
Cabbage	40	45-95	100
Cucumber	60	65-95	105
Eggplant	60	75-85	95
Pea	40	65-75	85
Pepper	60	65-75	85
Tomato	50	65-85	95

Optimum temperatures vary among crops

See UC Davis Extension: http://sacmg.ucanr.edu/files/164220.pdf



Fertilization

- Some media have trace nutrients
- After first or second set of true leaves, apply 1/4 strength liquid fertilizer weekly
- Well balanced N-P-K

Liquid synthetic

MircaleGro, Peters, Vigoro

Organic

- Fish emulsion (stinks!)
- Compost tea
- Rinse off any fertilizer that contacts foliage





Hardening-off Seedlings

Kicking the kids out of the house

1-2 weeks prior to transplant time, gradually expose to daytime outdoor conditions

- Protected from wind
- Shaded
- Reduce watering
- Bring in at night

Exceptions

- Harden-off tomatoes by reducing water
- Cucurbits and cauliflower very gradually





Night temps < 54°F affects fruit development



Setting Out

Planting transplants in garden

- Plants are ready to set out when their roots have filled the container and have several sets of true leaves
- Monitor watering closely check daily
- Mix in slow release or organic fertilizer at planting time, continue to liquid feed for few weeks



Healthy roots are white and firm

NC STATE EXTENSION

Extension Gardener


Plant Toolbox



https://plants.ces.ncsu.edu/

Select


'Find a Plant'



NC State Extension Homegrown

https://homegrown.extension.ncsu.edu

- In the Garden Videos
- In the Kitchen Videos
- On the Farm Videos





Need Help with Garden Problems?



Master Gardener | Chatham County

Plant Clinic: MW 1:00-4:00, F 9:00-12:00

chathamemgv@gmail.com

919-545-2715

(Except during COVID-19, email is preferred)



Send us your problems!

Questions we may ask:

- Crop and cultivar
- Describe signs and symptoms
 - Include photos!
- When you started noticing problems
- Cultural conditions
 - Light, soil, water, planting time etc.






Send Us Good Photos!

Photos should:

- Include healthy and unhealthy parts
- Have a scale object
- Be in focus
- Show an up-close image
- Show the whole plant
- The more, the better



Diagnosis: cataracts?



Subscribe to the

Chatham Gardener Newsletter

Chatham Gardener email list

- Sustainable gardening information
- Monthly email updates
- What to plant, pest alerts, timely tips
- Upcoming classes and events

To subscribe: https://chatham.ces.ncsu.edu/email-me/







Thank you!

matt_jones@ncsu.edu



